

Preparation and absorption spectra of aminothiazole derivatives for drug development studies

Dr. Y. Liu¹, Dr. H. Zhao¹, Dr. X. Chen¹, Dr. Q. Sun^{1*}

¹ School of Public Health and Preventive Medicine, Nanjing Medical University, Nanjing, China

ABSTRACT


A series of 4-aryl-2-aminothiazole derivatives were synthesized with an objective to develop novel and potent coupling components of synthetic origin as well as in dye synthesis. The required derivatives of 4-phenylsubstituted-2-aminothiazole were synthesized via a multicomponent condensation between thiourea, acetophenone and iodine. The intermediates were obtained using acetophenone and various substitute aldehydes to synthesize the intermediates which on cyclization with sulphur yielded the final products. Synthesized compounds were purified, characterized and evaluated for their spectra properties. All the synthesized intermediates exhibited moderate to significant properties. They were found to possess good coupling properties as well as high degree of brightness and a colour deepening effect compared to other heterocyclic couplers obtained from aniline and anthraquinone dyes.

Keywords: 4-phenyl-2-aminothiazole, synthesis, acetophenone, heterocyclic and condensation.

I. INTRODUCTION

Thiazole nucleus has been established as the potential entity in the largely growing chemical world of heterocyclic compounds possessing promising characteristics. Substituted thiazoles and their biheterocycles have received considerable attention during last two decades as they are endowed with wide range of therapeutic properties (Ananthanarayan, 2006). A number of thiazole derivatives have been reported to possess significant and diverse biological activities such as antimicrobial [Koti, 2006], analgesic, anti-inflammatory, antioxidant [Singh, 2011] and anti-allergic activities (Ishaq et al., 2012). Various pesticides possessing a thiazole nucleus are well known in agriculture. Large numbers of thiazole derivatives have emerged as active pharmaceutical ingredients in several drugs for their potential anti-inflammatory (Baur, 1996), anti-tumour (Azam, 2007), anti-hyperlipidemic, anti-hypertensive and several other biological properties. Besides, thiazoles are also synthetic intermediates and common substructures in numerous biologically active compounds (Cruikshank, 1975). Thus, the thiazole nucleus has been much studied in the field of organic and medicinal chemistry (Aoyama, 2006).

In continuation to these efforts and with an objective to develop potent heterocyclic intermediates of synthetic origin, it was decided to synthesize certain amino thiazole derivatives and evaluate them for their coupling potential in dye synthesis. The synthesis from acetophenone, thiourea and iodine as well as the absorption spectra of these intermediates are discussed.



2-amino-4-arylthiazole

II. MATERIALS AND METHOD

All the chemicals for the synthesis were purchased from approved vendors of different make like Aldrich, Dayang chemicals, Ievakluga, Synquest laboratory and King scientific and all chemicals were of laboratory grade. Completion of reaction were confirmed using physical constant determination (Sharp or narrow melting ranges). Melting points were determined using melting point apparatus in open capillaries and are uncorrected.

Further the compounds synthesized were proceeded for TLC wherein single spots (1st TLC run) were observed, indicating completion of reaction. After work-up was completed, (unreacted starting materials were removed by washing with ether), the products were subjected to purification by recrystallization process. Again TLC was run to find out exact R_f value. TLC plates used for final recrystallized product were pre-coated silica gel G plates. Solvent systems were developed using trial and error method by use of appropriate solvents of different polarity until the use of diethylether : ethanol (75%: 25%) solvent mixture was established. The Visible absorption spectra were measured using CARY 630 UV-visible spectrophotometer. Model: agilent technology . The infra-red spectra were carried out on FTIR Nexus 670 spectrophotometer in KBr disc (thermo nicolet) and absorption bands are expressed in cm⁻¹.

III. SYNTHESIS OF 2-AMINO-4-PHENYL THIAZOLE DERIVATIVES AII

3.1 Preparation of the 2- amino-4-phenylthiazole AII₁

Resublimed iodine (7.6g,0.03mol) was added to the slurry of the corresponding ketone (acetophenone and its para substituents- chloro, 2,4-dichloro, 3,4-dibromo, methyl, and bromo)(0.03mol) and thiourea (4.5g, 0.06mol) in toluene and the mixture was heated in an oil bath at 130°C - 140°C overnight. After cooling, the reaction mixture was diluted with distilled water (50ml) and heated to dissolve most of the solid, again cooled to ambient temperature. The aqueous solution was treated with 25% aqueous ammonium hydroxide(to pH 9-10). The precipitated thiazole was filtered, washed successively with water, collected and purified by crystallization from hot ethanol (Kamaljit et al, 2002).

3.2 Preparation of the 2- amino,4-(p-chlorophenyl) thiazole AII₂

The procedure is the same as in 3.1 above, except that the resublime iodine and thiourea were reacted with para-chloroacetophenone in toluene. The mixture was heated in an oil bath at 130°C - 140°C overnight. After cooling, the reaction mixture was diluted with distilled water (50ml) and heated to dissolve most of the solid, again cooled to ambient temperature. The aqueous solution containing the hydroiodide was treated with 25% aqueous ammonium hydroxide (to pH 9-10). The precipitated thiazole was filtered, washed successively with water, collected and purified by crystallization from aqueous ethanol (Kamaljit et al, 2002).

3.3 Preparation of the 2- amino,4-(p-aminophenyl) thiazole derivative AII₃

The procedure is the same as in 3.1 above, except that the resublime iodine and thiourea were reacted with para-aminoacetophenone in toluene and the mixture was heated in an oil bath at 130°C - 140°C overnight.

3.4 Preparation of the 2- amino-4-p(2,4-dichlorophenyl) thiazole derivative AII₄

The procedure is the same as in 3.1 above, except that the resublime iodine and thiourea were reacted with 2,4-dichloroacetophenone in toluene. The mixture was heated in an oil bath at 130°C - 140°C overnight.

3.5 Preparation of the 2- amino-4-(p-bromophenyl) thiazole derivative AII₅

The procedure is the same as in 3.1 above, except that the resublime iodine and thiourea were reacted with para-bromoacetophenone in toluene. and the mixture was heated in an oil bath at 130°C - 140°C overnight.

3.6 Preparation of the 2- amino-4-p(2,4-dibromophenyl) thiazole derivative AII₆

The procedure is the same as in 3.1 above, except that the resublime iodine and thiourea were reacted with 2,4-dibromoacetophenone in toluene. The mixture was heated in an oil bath at 130°C - 140°C overnight.

Scheme 1 ; Synthesis of 2-amino-4p(phenyl)thiazole derivatives

3.7 Recrystallization

All aminothiazole derivatives were purified by 3- 4 recrystallizations from hot ethanol. A known weight of the intermediate was dissolved in small quantity of ethanol and heated up. It was then filtered off using a Buchner funnel with a suction pump. The crystals were collected, washed severally with water and dried. After the recrystallization, the purity of each compound was checked by spotting on a thin layer chromatography plate

IV. RESULTS AND DISCUSSION

In a mixture of acetophenone and thiourea in toluene, iodine was added dropwisely with shaking. The mixture which was heated on an oil bath overnight at 130-140°C afforded 2-amino-4-aryl-thiazole(1). Compound 1 on reaction with 4-chloroacetophenone give 1-{4-[(4-phenyl-1,3-thiazol-2-yl) amino] phenyl}ethanone (2) which on further

stepwise reaction with various aromatic aldehydes afforded various 2-amino-4p-phenylthiazole derivatives respectively. The primary structural difference within the series involves the nature of various substituted aldehydes. Synthesized compounds were found to be crystalline in nature and easily soluble in ethyl acetate, benzene, DMSO and DMF but insoluble in hexane and toluene.

With the help of analytical techniques such as melting point, UV-visible spectrophotometer and FTIR, the synthesized derivatives were characterized. These compounds showed absorption band for C-S stretching of thiazole ring between 674-745 cm^{-1} and 1028-1088 cm^{-1} for C-N. All the compounds showed absorption peaks for different kinds of functional groups at their respective regions. All of them were found to be in full consonance with assigned structures.

For Compound AII₁: Yellow crystals were obtained; It was purified by recrystallization in hot ethanol. Melting point 144-147°C, 74% yield which correspond to literature values 148°C, 80% yield (Prajapati, 2010). FTIR (KBr)/ cm^{-1} : 1624 (C=C), 847 (CH-Ar bend), 2283 (Ar C-H), 3435 (NH str), 1483, 1532 (Ar C=C str), 1340 (C-C), 1039 (C-N), 691 (C-S), 3134 (Ar-H)

FOR COMPOUND AII₂: Off white crystals were obtained. It was purified by recrystallization in hot ethanol. Melting point 173°C, 67% yield 85% yield. FTIR (KBr)/ cm^{-1} : 1621 (C=C), 827, 741 (CH-Ar bend, double), 2847 (Ar C-H), 3391 (NH str), 1569 (Ar C=C str), 1088 (C-N), 1196 (C-H), 741 (C-S), 827 (C-Cl), 3119 (Ar-H).

FOR COMPOUND AII₃: A deep YELLOW crystal was obtained. It was purified by recrystallization in hot ethanol. Melting point 187-188°C, 52% yield. FTIR (KBr)/ cm^{-1} : 1610 (C=C), 831, 738 (CH-Ar bend, double), 2776 (Ar C-H), 3410 (NH), 1490, 1427 (Ar C=C str), 1265 (C-C), 1043 (C-N), 685 (C-S), 3116 (Ar-H), 1539 (N-H Ar)

FOR COMPOUND AII₄: A brownish yellow crystal was obtained, It was purified by recrystallization in hot ethanol. Melting point 190-192°C, 59% yield. FTIR (KBr)/ cm^{-1} : 1630, 1684 (C=C Ar), 827, 764 (CH-Ar bend, double), 2922 (Ar C-H), 3309 (NH), 1599 (NH Bnd), 1520, 1580 (Ar C=C str), 1632 (C-C), 784, 723 (C-Cl), 1028 (C-N), 674 (C-S), 3116 (Ar-H), 2922 (C-H Ar), 1129, 961 (C-H)

FOR COMPOUND AII₅: A BROWNISH YELLOW crystal was obtained, It was purified by recrystallization in hot ethanol. Melting point 185-187°C, 61% yield. FTIR (KBr)/ cm^{-1} : 1602 (C=C), 849, 771 (CH-Ar bend, double), 3432, (NH str), 1599 (NH Bnd), 1159 (C=C Ar), 1028, 1326 (C-C), 1086 (C-N), 693 (C-S), 693 (C-Br), 3060 (Ar-H).

FOR COMPOUND AII₆: A pale yellow crystal was obtained, It was purified by recrystallization in hot ethanol. Melting point 199-202°C, 67% yield. FTIR (KBr)/ cm^{-1} : 1621 (C=C Ar), 823 (CH-Ar bend, double), 2847, 2948 (Ar C-H), 3384 (NH str), 1565 (NH Bnd), 1490 (C=C Ar str), 1319 (C-C), 1069 (C-N), 741 (C-S), 741 (C-Br), 1189 (C-H Ar), 3116 (Ar-H).

V. CONCLUSION

The analytical and other informational data, available in literature so far, have rendered thiazole significantly important class of heterocyclic compounds and their applications in ever challenging dye synthesis of various classifications. All the synthesized intermediates exhibited moderate to significant properties. They were found to possess good coupling properties as well as high degree of brightness and a colour deepening effect compared to other heterocyclic couplers obtained from aniline and anthraquinone dyes.

This particular research study, in reference, would extend great deal of help to researchers in reckoning and determining the best and most productive, economical, suggestive and conclusive access to various thiazoles of organic and synthetic importance superseding other coupling compounds of their class.

REFERENCES

1. Ananthanarayan R. And Jayaram Paniker (2006). TextBook of Microbiology; Orient Blackswan; 5th Edition; 39
2. Aoyama T., Murata S., I. Arai, N. Araki, T. Takido, Y. Suzuki, M. Kodomari (2006) Tetrahedron; 62; Pp 3201-3213.

3. Azam .F. et.al (2007). Synthesized Schiff bases of naphtha [1, 2-d] thiazol-2-amine and metal complex of 2-(2'hydroxy) benzyldineaminonaphthothiazole, J Zhejiang Univ Sci. B ;8(6), pp 446-452
4. Baur, R. W, Kirby M. D and Sherris J. C. (1996). American Journal of Clinical Pathology. Pp 45, 493-496.
5. Cruikshank R., Duguid J. P ,Marmion B. P and . Swain R. (1975). Medical Microbiology; Churchill Livingstone, New York; 2nd Edition; 90.
6. IshaTomer, Raghav Mishra Priyanka, N. K. and Sharma, K. K. (2012). Synthesis and antimicrobial evaluation of some novel thiazole derivatives .Pelagia Research Library, 2012, 3 (3) : 361-366
7. Lu Y et.al (2009). Synthesis of 4 -substituted methoxybenzoyl-aryl-thiazole Journal of Medical Chemistry.52(6); 1701-11
8. Patel K. H; A. G. (2006). Mehta synthesized Thiazolidone derivatives of 2-Amino-6-(2- naphthenyl) thiazolo [3,2-d] . E-Journal of Chemistry; 2006 ;3(13), pp267-273.
9. Pattan S. R. et al (2009). Synthesis of amino derivatives of thiazole. Asian Journal of Research Chemistry. Pp2;2
10. Petrikaite.et.al (2007). Synthesis of thiazolidone-4-sulfmethizole-2-substituent derivatives,.Biologija.Pp53 ; 3, 45-50.
11. Prajapati A.K.,Vishal modi. Synthesis and biological evaluation of some substituted aminothiazole derivatives. J.chil chemical society 55NoC PP 240-243.